

30.07.04



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

Bescheinigung

Certificate

Attestation

REC 01 OCT 2004

WIPO

PCT

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03017323.1

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

R C van Dijk



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

PCT/EP200 4 / 0 0 8 5 8 7

30.07.03

Anmeldung Nr:  
Application no.: 03017323.1  
Demande no:

Anmeldetag:  
Date of filing: 31.07.03  
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Emdoka bvba,  
Drug registration and Marketing  
J. Lijsenstraat 16  
2321 Hoogstraten  
BELGIQUE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:  
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.  
If no title is shown please refer to the description.  
Si aucun titre n'est indiqué se referer à la description.)

Veterinary aqueous injectable suspensions containing florfenicol

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)  
revendiquée(s)  
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/  
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of  
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL  
PT RO SE SI SK TR LI

31. Juli 2003

EMDB0003 (933/03)  
D67/D894 KF/KF-ch

### Veterinary Aqueous Injectable Suspensions Containing Florfenicol

The invention relates to pharmaceutical compositions for veterinary use containing Florfenicol as the active pharmaceutical ingredient in high concentration, more specifically in aqueous injectable suspensions.

Florfenicol [D-(threo)-1-p-methylsulfonyl phenyl-2-dichloroacetamido-3-fluoro-1-propanol] is a known antibacterial agent which is commonly used for veterinary purposes. It is a broad-spectrum antibiotic that is active against many pathogenic veterinary bacteria (NEU et al., Antimicrob. Agents Chemother., 1980, 18, 2, 311-316; SYRIOPOULOU et al., Antimicrob. Agents Chemother., 1981, 19, 2, 294-297).

Injectable formulations containing Florfenicol are already described in several patents.

European Patent 0 014 437 B1 discloses not only the way of synthesis of Florfenicol, but also different formulation examples, including oral suspensions, injectable solutions in a water/dimethylacetamide mixture and sterile powders for extemporaneous reconstitution.

European Patent 0 546 018 B1 discloses a highly concentrated injectable formula for veterinary use. This injectable formula is based on a water-free combination of two different organic solvents. An additional viscosity reducing agent is required to enable the parenteral administration of this highly concentrated solution.

Although this formula is claimed to exhibit acceptable tissue tolerance at the injection site, practicing veterinarians designate its injection as very painful.

This injectable formula of European Patent 0 546 018 B1 might possibly be an improvement over former injectable solutions, e.g. formulations dissolved in pure N-Methyl-2-pyrrolidone, it still remains very irritating and painful, if compared to other physiologically acceptable aqueous solutions and suspensions, containing lower concentrations of physiologically acceptable solvents, with physiologically acceptable pH-values and distinctly less osmotic stress at the injection-site.

It is an object of the present invention to provide, highly concentrated aqueous injectable suspensions of the poorly water-soluble Florfenicol that can be manufactured with very good shelf life and stability, with a distinctly better local tolerance than the above mentioned injectable solutions, with short or sustained release serum levels for a long lasting antimicrobial activity after a reduced number of injections, and with limited sedimentation on standing, quick resuspendibility on gentle shaking, even after shipping, and easy syringibility on extraction.

Accordingly, the present invention relates to pharmaceutical compositions for veterinary use in the form of an aqueous suspension comprising Florfenicol or Florfenicol in the form of a substantially water-insoluble complex, co-crystal or salt, sterile and micronised, in a concentration up to 500 mg/ml.

With Florfenicol, such suspensions have not been available, so far.

In this context it shall be mentioned, that aqueous should be understood as essentially pure aqueous solutions without any further addition of organic solvents.

However it is evident, that these kinds of suspensions are not confined to injectable compositions. A further application in the form of stable, highly concentrated suspensions of Florfenicol, to be diluted into drinking water and animal-milk and/or milk replacers for veterinary use, appear also feasible.

In the following description of these formulations, all the concentrations of the ingredients mentioned are expressed in mg/ml.

Accordingly, the active ingredient Florfenicol is used in these aqueous suspensions in concentrations up to 500 mg/ml.

The Florfenicol raw material has to be sterile.

The Florfenicol raw material has to be micronised also, i.e.  $\geq 90\%$  of the total volume of the Florfenicol raw material consists of particles with a particle size between 0.5  $\mu\text{m}$  and 200  $\mu\text{m}$ , preferably between 1  $\mu\text{m}$  and 100  $\mu\text{m}$ , more preferably between 1  $\mu\text{m}$  and 75  $\mu\text{m}$  and most preferably between 1  $\mu\text{m}$  and 50  $\mu\text{m}$ .

The grade of micronisation determines the shape of the Florfenicol serum curve in treated animals, viz.: the smaller the particles are, the faster the parenteral resorption occurs, but also the higher the tendency is of the suspension towards the detrimental "Ostwald ripening". Smaller particle sizes are used for treatments where high initial serum levels are required. Bigger particles are used for longer lasting serum levels. An equilibrated combination of both produces a serum curve with a bactericidal initial serum peak and a long lasting bacteriostatic serum slope for a long antimicrobial protection. So, the assortment of a well-equilibrated range out of this particle size distribution is a good method to provide a particular retardation of the active pharmaceutical ingredient, if required.

Micronisation of the particles should be conducted according to the state of the art, in dry form by means of a jet mill or in suspended form by means of a Co-Ball mill. The particles should preferably be non-acicular, but tabular for optimal resuspendibility.

Well resuspendable aqueous suspensions contain a series of necessary excipients, viz.: buffers, suspending agents, surface-active agents, antioxidants and antimicrobial preservatives. Some of these constituents do flocculate the suspension, whereas others deflocculate them. The optimal equilibrium between flocculation and deflocculation has been established experimentally, with the

aim to obtain suspensions with limited sedimentation on standing, quick resuspendibility on gentle shaking, even after shipping, and easy syringibility on extraction.

For an optimal chemical and physical stability of the suspension, the bulk solution needs to be buffered. Buffering components can be, but are not limited to: citric acid, ascorbic acid, isoascorbic acid, maleic acid, tartaric acid and boric acid and their respective hydrosoluble salts, monobasic and/or dibasic hydrogen phosphate, basic or acidic amino acids, such as arginine, cysteine, etc. and their hydrosoluble salts, as well as mixtures thereof. These buffer mixtures are present in concentrations from 1 mg/ml to 250 mg/ml, more preferably between 50 mg/ml and 250 mg/ml. It was observed that adequately buffering the injectable suspension, at the same time dramatically improves the local tolerance at the injection-site.

The pH of the bulk has to be within the range of physiological acceptance, which means between 3 and 9, more preferably between 5 to 8 and most preferably between 5 and 7.

The pH-value for optimal stability of Florfenicol in aqueous solution lies between 3 and 6. However, Florfenicol has a low solubility in water: at pH = 7.0 it dissolves into water in a concentration of not more 1.3 mg/ml only, whereas Chloramphenicol dissolves in a concentration of 2.5 mg/ml.

Different suspending agents are used.

To equilibrate the floating density of the continuous phase and the repellent forces between adjacent particles, a sugar such as maltosé, fructose and glucose but not limited thereto, a polyhydric alcohol such as sorbitol, mannitol and xylitol but not limited thereto, sugar acids and/or fruit acids with at least 3 functional hydroxy and/or carboxy groups such as gluconic acid, heptagluconic acid, saccharic acid and citric acid but not limited thereto, and their water-soluble alkaline metal salts such as lithium, sodium, potassium, earth-alkaline metal salts such as magnesium and calcium salts and physiologically acceptable transition metal salts such as iron, manganese, zinc as well as their

double salts can be added. Their concentration amounts from 10 mg/ml to 400 mg/ml, more preferably between 50 mg/ml and 300 mg/ml.

Dissolved in water, Florfenicol readily hydrolyses. Hence the first challenge of the galenical development of these Florfenicol suspensions consisted in keeping the suspended material as much as possible out of dissolution, by repelling its dissolution through the addition of sufficient amounts of readily hydrosoluble excipients. Increasing the concentrations of both the buffer mixtures and the suspending agents in the continuous phase noticeably reduced the degradation of Florfenicol.

For an optimal homogenisation of the floccules, sodium carboxymethylcellulose (Na-CMC) of low or medium viscosity grade can be added. It is used in a concentration of 0.1 mg/ml to 10 mg/ml. It increases the electrical charge of the particles in suspensions, up to its maximum value, which is mostly reached around a concentration of 5 mg/ml (POLDERMAN, Am. J. Hosp. Pharm., 1962, 19, 611-620).

Polyvinylpyrrolidone (PVP) is a non-ionizable compound that renders the suspension a better local tolerance at the injection-site. It however can enhance caking. To avoid caking and to improve resuspendibility, the content of polyvinylpyrrolidone should be in equilibrium with the content of the sodium carboxymethylcellulose added. The amount of polyvinylpyrrolidone should be between 0.3 and 30 mg/ml, preferably between 1 and 10 mg/ml. Different polyvinylpyrrolidones with K-values between K 12 and K 32 can be used, but the ones with the lowest molecular weight are physiologically the most acceptable. In this way, polyvinylpyrrolidones with K-values from K 12 to K 15 are preferred.

The presence of a surface-active agent is necessary for two reasons: as a wetting agent for the particles to be suspended and for diminishing the risk of "Ostwald ripening", viz.: to suppress crystal growth during shelf life. These agents can be, but are not limited to: choline, phospholipids such as lecithine, non-ionogenic surface active agents, such as poloxamers, sorbitan fatty acid esters and polyoxyethylene sorbitan fatty acid esters, known under several commercial names, e.g.: spans and tweens, arlacels, crills and montanes. Lecithins are the preferred surface-active agents in the present suspensions.

They consist of mixtures of phospholipids with different purity grades. The lecithins can be used in two different ways: either coated on the Florfenicol particles, in concentrations varying from 0.1 to 20 mg/ml, or dispersed into the continuous phase in concentrations from 0.1 to 50 mg/ml. Useful for that purpose are e.g. Lecithin brands Phospholipon® 90G, 90H and 100H from Nattermann Phospholipids GmbH, Köln. Other non-ionogenic surface-active agents can be used together with or instead of lecithins, e.g.: sorbitan esters or polyoxyethylene sorbitan esters in concentrations of 1 to 30 mg/ml.

Suspensions have to contain one or more water-soluble antioxidants and respective synergists. Useful antioxidants are sodium formaldehyde sulfoxylate, sodium metabisulphite, sodium phosphite, ascorbic acid, tartaric acid and cysteine, but not limited thereto, used alone or in combination, in concentrations of 0.1 to 40 mg/ml. Synergists potentiate the activity of these antioxidants. Useful synergists are: chelating or sequestering agents, such as citric and tartaric acid and especially sodium edetate and other edetic acid derivatives or the like. They are used in concentrations from 0.01 to 5 mg/ml.

Finally, the multi-dose injectable suspensions should be protected against microorganisms, possibly invading with contaminated injection needles, and hence have to meet the "Test for efficiency of antimicrobial preservation", adopted in the European Pharmacopoeia 4 "under 5.1.3 – 1 Parenterals". With that aim, they should contain one or more antimicrobial preservatives. Adequate and physiologically acceptable preservatives are, e.g. the parabens (methylparahydroxybenzoate, ethylparahydroxybenzoate, etc.), benzyl alcohol, chlorbutanol, phenoxyethanol, phenol and derivatives, quaternary ammonium compounds such as benzalkonium chloride and benzethonium chloride, benzoic acid, sorbic acid and their alkaline salts.

The suspensions of the present invention meet all the characteristics of good and reliable injectable suspensions, viz.: they show a limited sedimentation on standing as well as after shipping, a good resuspendibility on gently shaking and a good syringibility, even with thin needles.

The sedimentation height of the suspensions is investigated according to the method of WARD & KAMMERMYER, Ind. Eng. Chem., 1940, 32, 5, 622-626. It

is expressed as the Hu/Ho-value, and amounts not less than 90% after 24 hours.

The resuspendibility of the suspensions is measured according to the method of MATTHEWS & RHODES, J. pharm. Sci., 1968, 57, 569-573. After normal standing at room-temperature, the complete resuspendability is achieved after less than 10 rotations. Moreover, after a "shipping test" of 2000 km the suspensions can also easily be resuspended on gentle shaking, and do not stick at the bottom of the vial.

The syringibility of the suspension is expressed by means of the time needed to completely fill a 10 ml syringe, equipped with a 40 mm needle with diameter of 21G (0.8 mm). At room-temperature, the syringibility-time is less than 20 seconds.

The suspensions should be filled in siliconised sterile glass vials or sterile polyethylene or polypropylene vials, conceived for parenteral use, closed with sterile butylrubber stoppers and alu caps.

The following example describes the invention in general:

## General Example

**Active pharmaceutical ingredient, duly micronised** up to 500 mg/ml

Buffer mixture with pH between 5 and 8

1 to 250 mg/ml

**Suspending agents:**

<p><b>Sugars, polyhydric alcohols, sugar acids, fruit acids and their salts for equilibrating the floating density of the continuous phase and the repellent forces between adjacent particles</b></p>	<p><b>10 to 400 mg/ml</b></p>
--	-------------------------------

**Sodium CMC, low or medium viscosity grade** 0.1 to 10 mg/ml  
preferably  $\leq 5$  mg/ml

**PVP** 0.3 to 30 mg/ml  
preferably 1 to 10 mg/ml

<b>Surface active agents</b>		
<b>or: phospholipids</b>	<b>coated on the particles</b>	<b>0.1 to 20 mg/ml</b>
	<b>dispersed in the continuous phase</b>	<b>0.1 to 50mg/ml</b>
<b>and/or: non-ionogenic surface active agents</b>		<b>1 to 30 mg/ml</b>

<b>Antioxidants</b>	<b>0.1 to 20 mg/ml</b>
---------------------	------------------------

<b>Antioxidant synergists</b>	0.01 to 3 mg/ml
-------------------------------	-----------------

## Antimicrobial preservatives

## Water for injection

The following Examples describe the invention in detail:

#### Example 1

Florfenicol micronised	300 mg
Sodium citrate	50.0 mg
Monopotassiumdihydrogenphosphate	1.5 mg
Glucose monohydrate	80.0 mg
Sodium CMC low viscosity	2.0 mg
Polyvinylpyrrolidone K12	3.5 mg
Lecithin	3.0 mg
Sodiumformaldehyde sulfoxylate	2.5 mg
Sodium metabisulphite	0.1 mg
Sodium edetate	0.1 mg
Methylparaben	1.0 mg
Propylparaben	0.5 mg
Water for injection q.s. ad	1 ml

#### Example 2

Florfenicol micronised	250 mg
Sodium citrate	50.0 mg
Sorbitol 70%	250.0 mg
Sodium CMC medium viscosity	0.8 mg
Polyvinylpyrrolidone K15	3.5 mg
Polysorbate 80	1.0 mg
Sorbitan monooleate	1.0 mg
Sodiumformaldehyde sulfoxylate	2.5 mg
Sodium phosphite	10.0 mg
Sodium edetate	0.1 mg
Benzyl alcohol	9.0 mg
Water for injection q.s. ad	1 ml

**Example 3**

Florfenicol micronised	400.0 mg
Sodium citrate	80.0 mg
Monopotassiumdihydrogenphosphate	10.0 mg
Sodium gluconate	150.0 mg
Sodium CMC low viscosity	1.5 mg
Polyvinylpyrrolidone K12	3.5 mg
Lecithin	3.0 mg
Sodiumformaldehydesulfoxylate	2.5 mg
Sodium metabisulphite	0.1 mg
Sodium edetate	0.1 mg
Methylparaben	1.0 mg
Propylparaben	0.5 mg
Water for injection q.s. ad	1 ml

31. Juli 2003

CLAIMS

1.           Pharmaceutical compositions for veterinary use in the form of an aqueous injectable suspension comprising Florfenicol or Florfenicol in the form of a substantially water-insoluble complex, co-crystal or salt, sterile and micronised, in a concentration up to 500 mg/ml.
2.           The compositions of claim 1, wherein the Florfenicol is present as sterile raw material, in a particle size with  $\geq 90\%$  of the total volume of the Florfenicol raw material consisting of particles with a particle size between 0.5  $\mu\text{m}$  and 200  $\mu\text{m}$ .
3.           The compositions of claim 1 or 2 wherein the continuous phase of the suspension contains 1 to 250 mg/ml of a buffer mixture to provide a pH-value in the range of 5 to 8.
4.           The composition of any of claims 1 to 3 wherein the suspension is stabilized by at least one sugar, polyhydric alcohol, sugar acid and/or a fruit acid with at least 3 functional hydroxy and/or carboxy groups, or a salt thereof in a concentration of 10 to 400 mg/ml.
5.           The compositions of any of claims 1 to 4, comprising sodium carboxymethylcellulose as a suspending agent in a concentration of 0.1 to 10 mg/ml.
6.           The compositions of any of claims 1 to 5, comprising at least one injectable grade polyvinylpyrrolidone in concentrations of 0.3 to 30 mg/ml.

7. The compositions of any of claims 1 to 6, wherein a phospholipid surface-active agent is incorporated in a concentration of 0.1 to 50 mg/ml, either coated on the particles or dispersed into the continuous phase and/or other non-ionogenic surface active agents, in a concentration of 1 to 30 mg/ml.
8. The compositions of any of claims 1 to 7, comprising common additives such as antioxidants and their synergists, and antimicrobial preservatives.
9. The compositions of any of claims 1 to 8, comprising Florfenicol as a sterile injectable suspension, filled aseptically in sterile primary packing materials.
10. The compositions of any of claims 1 to 9 for therapeutic and prophylactic use in animals.

**ABSTRACT**

This invention relates to pharmaceutical compositions for veterinary use in form of aqueous injectable suspensions comprising Florfenicol or Florfenicol in form of a substantially water-insoluble complex, co-crystal or salt, sterile and micronised, in a concentration up to 500 mg/ml. The suspensions enable parenteral antimicrobial therapies in animals, with limited numbers of injections and showing good general- and local tolerances. They also possess a limited sedimentation on standing as well as after shipping, are easily resuspendable and have good syringibility.